

"Stobaeana", named after the founder of the Museum, Kilian Stobaeus, (1690–1742), contains contributions – based on the collections housed in the museum – within the fields of systematics, taxonomy, biogeography etc. Also reports on ongoing projects associated with the Museum might be considered for publication.



## A new species of *Acrosathe* (Diptera, Therevidae) from the Baltic Islands

HUGO ANDERSSON

Andersson, H.: A new species of *Acrosathe* (Diptera: Therevidae) from the Baltic Islands. *Stobaeana* 1994(2):1–6. Lund, Sweden 6 November 1994. ISSN 1104-5957.

A new species, *Acrosathe baltica* n. sp., from the Baltic Islands is described and figured. It is compared with the closely related species *A. sybarita* Loew from south and southeast Europe and the in the area common species *A. annulata* Fabricius.

H. Andersson, Swedish Mus. Nat. Hist., c/o Zoological Dept., Helgonavägen 3, S-223 62 Lund, Sweden.

In 1986, Lyneborg revised the Old World species of the genus *Acrosathe* Irwin & Lyneborg, 1981. Of the nine species treated four had an eastern palaearctic distribution, found in The Far East Territory of the U. S. S. R., China, Japan and Taiwan, one was known only from West Siberia and Kazakhstan, while two had a southern European distribution, South Spain resp. Italy and the South European Territory of the U. S. S. R. Only one species, *A. annulata* Fabricius, was found to have a wide European distribution. In 1988, Nagatomi & Lyneborg described further two species from Japan.

*A. annulata* was divided into four geographical races supposed to have developed in different Mediterranean refugia during the pleistocene glaciations (Lyneborg 1986:112). The race (form) A was found in Denmark, Fennoscandia, East and Central Europe south to the black sea, Rumania, North Greece and Northwest Italy. In Sweden *annulata* form A has been recorded from the provinces Sk, Bl, Ha, Sm, Ö, G, Bo, Dr, Ån, Nb, and Ly. It has been regarded as a species restricted to sandy coastal localities. Ardö (1957:168) found it bound to the marine shore dune ecosystem. In Sweden it is found also in a number of sandy inland localities, some of which are

clearly connected to old sea shores, developed during various periods of the deglaciation era. In 1987, Mikael Sörensson (Sörensson 1989) made a survey of the insect fauna of the sand dune area Ulla Hau in the small island Fårön, at the north end of the Baltic island Gotland. I was entrusted the determination of the Diptera from this investigation and found to my surprise that the material of *Acrosathe* did not, as expected, belong to *annulata* but to a species closer related to *A. sybarita* Loew. It will be described as new below.


### *Acrosathe baltica* sp. n.

Figs. 2, 4, 5, 7, 9, 12, 15, 18, 22–24.


**Type material:** *Holotype*: ♂, Sweden, Gotland, Gotska Sandön, 1922–23, A. Jansson. — *Paratypes*: 1♂ 3♀, same data as holotype; 2♂ 1♀, Gotland, Fårö, Ulla Hau, 3–16.VI.1987 (1♂ 1♀), 16.VI–27.VIII.1987 (1♂), M. Sörensson. Types in Zoological Museum, Lund.

**Etymology:** The name derives from the type localities: islands in the Baltic Sea.


**Diagnosis:** *Acrosathe baltica* n. sp. is closely related to *A. sybarita* Loew. It differs from *sybarita* by a smooth, not dentated dorsal



Figs. 1-8. — 1, 3, 6. *Acrosathe annulata* Fabr. (SE: Sk. Sandhammaren). — 2, 4, 5, 7. *A. baltica* n. sp. (Paratype: S: Go. Ulla Hau). — 8. *A. sybarita* Loew. (I: Firenze, Cascine). — 1-2. Epandrium and cerci, dorsal view. — 3-4. Male sternite 8, ventral view. — 5. Hypandrium, ventral view. — 6-7. Gonocoxites with appendages, internal view. Most of the bristles on gonocoxa omitted. — 8. Appendages of gonocoxa. — GC - gonocoxa; GS - gonostyle; PP - parameral process; VL - ventrale lobe.



Figs. 9–18. — 9, 12, 15, 18. *Acrosathe baltica* n. sp. (Paratype: SE: Go. Ulla Hau). — 10, 13, 16. *A. sybarita* Loew. (I: Firenze, Cascine). — 11, 14, 17. *A. annulata* Fabr. (S: Sk. Sandhammaren). — 9–10. Gonocoxa resp. apex of gonocoxa with appendages, external view, most of the bristles of the gonocoxa omitted. — 11–13. Aedeagus, lateral view. — 14–15. Aedeagus, ventral view. — 16. Aedeagus, dorsal view. — 17. Digging spines of female tergite 10. — DAP - dorsal aedeagal apodeme; EAP - ejaculatory apodeme; PLP - posterolateral process of aedeagus; VAP - ventral aedeagal apodeme. - A - posterodorsal digging spines; B - anterolateral digging spines.



Figs. 19–27. — 19–21. *Acrosathe annulata* Fabr. (SE: Sk. Sandhammaren). — 22–24. *A. baltica* n. sp. (Paratype: S: Go. Ulla Hau). — 25–27. *A. sybarita* Loew. (I: Pisa, Tirvenia). — 19, 22, 25. Female tergite 8. — 20, 23, 26. Vaginal apodem. — 21, 24, 27. Female sternite 8.

aedeagal apodeme (Fig. 12, cf Fig. 13), by the gonostylus distinctly broadened at middle, by parameral process thicker and not so sharply pointed at apex and by ventral lobe smooth, not granulated on the dorsal margin (Fig. 7 cf. Fig. 8). The female of *baltica* differs from *sybarita* and *annulata* in having the small erect hairs on abdominal segments 6–8 white instead of black. The upper frons and the vertex are distinctly brown as in *annulata*, not grey as in *sybarita*. The male of *baltica* differs from *annulata* in the sharply pointed posterolateral processes of aedeagus (Fig. 12, cf Fig. 11) and the rounded posteriolateral margins of the epandrium (Fig. 2, cf Fig. 1).

### Description

The new species *Acrosathe baltica* fit the description of the genus (Lyneborg 1986:102). In its general features it is similar to *annulata*. In the following description only characteristics by which it is similar or dissimilar to *annulata* and/or *sybarita* will be treated.

Male. — Frons above the antennae with shorter, proclinate, white pile as in *annulata*; in *sybarita* this pile is longer, erect and partly (behind) black.

Mesonotum with white pile as in *annulata*; in *sybarita* the white pile is mixed with black hairs, especially on the central part of the anterior half.

Anterior femora with usually 1 (0–2) *av*; in *annulata* 2 (0–3) *av* and in *sybarita* 2 *av*. Middle femora without *av*; in *annulata* usually 2 (1–2) and in *sybarita* 2 (?) *av*. Hind femora with 4–5 *av*; in *annulata* and *sybarita* 5–6 *av*.

Epandrium with rounded posterolateral margins (Fig. 2); in *annulata* they are square (Fig. 1) and in *sybarita* not quite so sharply rounded. These differences in the epandrium can usually be seen without preparation. The ventral lobe (Fig. 7 VL) is slender with smooth dorsal margin and without vestiture of small microchaetae. In *annulata* and *sybarita* VL is shorter and broader with rugulose dorsal margin and with microchaetae on the internal (*annulata*, Fig. 6) or ventral (*sybarita*, Fig. 8) surface. The gonostyle (Fig. 7 GS) is very broad at middle; in *annulata* and *sybarita* distinctly slender. The parameral process (Fig. 7 PP) is rather thick and apically blunt; in *annulata* and *sybarita* PP is slender and in *sybarita* very acute at apex. The posterolateral processes of aedeagus (Figs. 12, 15 PLP) are slender and sharply pointed as in *sybarita* (Figs. 13, 16). In *annulata* PLP are blunt and with a dorsal triangular extension (Figs. 11, 14). The dorsal aedeagal apodeme (Figs. 12, 15 DAP) is slender without any dentations as in *sybarita* (Figs. 13, 16). The ventral aedeagal apodeme (Figs. 12, 15 VAP) is smaller and short. The ejaculatory apodeme (Fig. 12 EAP) is short and rather straight; in *sybarita* (Fig. 13) it is longer and apically upcurved.

Female. — Anterior part of frons has grey toment and white pile; the posterior part is chocolate brown with

short, black setae as in *annulata*. In my specimen of *sybarita* the whole frons is grey with only very weak indications of a brownish band at middle.

Mesonotum with adpressed white pile and longer, erect white hairs some of which are brownish. In *annulata* the adpressed pile is white but the longer, erect hairs are brown to black all over. The adpressed pile in *sybarita* is white at sides but brown along a middle stripe and the erect hairs are brown.

Middle femora have no *av*; in *annulata* there are 2–3 *av*.

The abdominal segments 1–7 have a thick, grey toment. On segments 2–4 there is an anterior area with black toment. On tergite 2 the black area reaches at middle about 2/3 of the length of the tergite; the black area is gradually narrowing towards the sides and does not reach the margins of the tergite. The black area of tergite 3 has about the same relative length but it reaches as a narrow brown band the lateral margins of the tergite. On tergite 4 the visible black area is about as broad as the grey area; it reaches the lateral margins of the tergite for about 1/4 of the length of the tergite. Tergites 5–7 are without significant darkening, possibly the extreme hind margins are somewhat brownish. Segment 8 is subshining with a very faint grey toment. In *annulata* the dark bands are narrower, about half the length of the visible part of the tergite. In *sybarita* the central part of tergite 1 is brownish. On tergites 2–3 the black spots are more circular and covers more than 3/4 of the length of the tergites. The spot of tergite 2 is halfcircular and laterally so restricted that it does not at all reach the downbent part of the tergite. The spot on tergite 3 reaches laterally the downbent parts of the tergite. The spot on tergite 4 is more transverse and narrowly reaches the sidemargins of the tergite. In *baltica* the grey parts of tergites 1–4 have an adpressed white pile but on the central disk of tergite 1 there is dark brown adpressed pile. The adpressed pile on the black spots of tergites 2–4 is dark brown. At the sides of T 1–2 there is a longer more or less erect white pile. T 5–8 and lateral parts of T 4 have numerous short erect white hairs. S 2–3 have longer white erect hairs while S 4–8 have short erect white hairs. The pile in *annulata* and *sybarita* is the same except that the short erect hairs on segment 4–8 are quite black. The posterodorsal digging spines of *baltica* on T 10, 6–7 on each side, are dark, moderately long and stout. The anterolateral digging spines on T 10 are about 4–5. weak, pale, slender and at most as long as the posterodorsal ones (Fig. 18). In *annulata* and *sybarita* the posterodorsal spines are relatively shorter and stouter, about 7, and the anterolateral ones are 6, strong, black and distinctly longer than the posterodorsal ones (Fig. 17). The vaginal apodeme of *baltica* (Fig. 23) is much shorter and broader than that of *annulata* and *sybarita* (Figs. 20, 26). Also T 8 and S 8

are shorter than in *annulata* and *sybarita* (Figs. 22, 24 cf Figs. 19, 21 resp. 25, 27).

#### Acknowledgments

The author is grateful to Mikael Sörensson, Lund, for the possibility to study the interesting Diptera-material from Fårön, to Roy Danielsson, Zoological Museum, Lund, for loan of material and to Teresa Mireström for technical assistants.

#### References

- Ardö, P. 1957. Studies on the marine shore dune ecosystem with special reference to the Dipterous fauna.— Opusc. ent. Suppl. XIV:1–255.
- Lyneborg, L. 1986. The genus *Acrosathe* Irwin & Lyneborg, 1981 in the Old World (Insecta, Diptera, Therevidae). — Steenstrupia 12(6):101–113.
- Nagatomi, A. & Lyneborg, L. 1988. The Japanese *Acrosathe* (Diptera, Therevidae). — Kontyu\* 56(3):600–617.
- Sörensson, M. 1989. Insektfaunan i Ulla Håu och några andra godländska sanddynområden. — Länsstyrelsen i Gotlands län, Naturvårdsfunktionen. Pp. 1–122.

# Stobæana

Occasional papers published by the Museum of Zoology, Lund University, Sweden

“*Stobæana*”, named after the founder of the Museum, Kilian Stobæus, (1696–1742), contains contributions — based on the collections housed in the museum — within the fields of systematics, taxonomy, biogeography etc. Also reports on ongoing projects associated with the Museum might be considered for publication.

Editor: Dr. Lennart Cederholm, Museum of Zoology, Lund University, Lund, Sweden.

“*Stobæana*” is published at irregular intervals and distributed on exchange basis to museums and other research institutions. Thus, we are anxious to get in touch with such institutes in the fields of zoological systematics, taxonomy, biogeography publishing papers/journals available through exchange.

Please contact:

Museum of Zoology, Helgonavägen 3, S-223 62, Lund, Sweden. Phone +46 (46) 10 93 30; Fax +46 (46) 10 45 41.

Additional copies can also be purchased at a cost of US \$5 per extra copy. Standing orders are accepted.

#### Stobæana

1. Scheller, U., Brinck, P. & Enckell, P. H. First record of Pauropoda (Myriapoda) on Borneo. 1994, 14 pp.
2. Andersson, H. A new species of *Acrosathe* (Diptera, Therevidae) from the Baltic Islands. 1994, 6 pp.
3. Nartshuk, E. P. New species and new records of Chloropidae from Afghanistan (Diptera). 1994, 5 pp.

1. Reports from the Lund University Ceylon Expedition in 1962. Vol. 1, 1971. XXXVI + 292 pp.....	33
2. Palm, T.: Die skandinavischen Elateriden-Larven (Coleoptera). 1972, 63 pp. ....	9
3. Nyholm, T.: Die nordamerikanischen Arten der Gattung <i>Cyphon</i> Paykull (Coleoptera). 1972. 100 pp. ...	11
4. Reports from the Lund University Ceylon Expedition in 1962. Vol. II. 1975. IV+ 239 pp. ....	30
5. Lindroth, C. H., Andersson, H., Böddvardsson, H. & Richter, S. H.: Surtsey, Iceland. The development of a new fauna, 1963—1970. Terrestrial invertebrates. 1973. 280 pp. ....	28
6. Birket-Smith, S. J. R.: On the abdominal morphology of Thysanura (Archaeognatha and Thysanura s.str.). 1974. 67 pp. ....	11
7. Möller Andersen, N.: The <i>Limnogonus</i> and <i>Neogerris</i> of the Old World with character analysis and a reclassification of the Gerrinae (Hemiptera: Gerridae). 1975. 96 pp. ....	14
8. Andersson, H.: Taxonomic and phylogenetic studies on Chloropidae (Diptera) with species reference to Old World genera. 1977. 200 pp. ....	24
9. Heie, O. E.: Revision of the aphid genus <i>Nasonovia</i> Mordvilko, including <i>Kakimia</i> Hottes & Frison, with keys and descriptions of the species of the world (Homoptera: Aphididae). 1979. 104 pp. ....	17
10. Sæther, O. A. (Ed.): Recent developments in chironomid studies (Diptera: Chironomidae). 1979. 150 pp. ....	25
11. Cederholm, L. (Ed.): Reports from the Lund University Ceylon Expedition in 1962. Vol. 3. 1982. 274 pp. ....	37
12. Huggert, L.: Revision of the West Palearctic species of the genus <i>Idris</i> Förster s. l. (Hymenoptera, Proctotrupoidea: Scelionidae). 1979. 60 pp. ....	10
13. Griffiths, G. C. D.: Studies on boreal Agromyzidae. XIV. <i>Chromatomyia</i> miners on Monocotyledones (Diptera: Agromyzidae). 1980. 61 pp. ....	11
14. Sæther, O. A.: Glossary of chironomid morphology terminology (Diptera: Chironomidae). 1980. 51 pp. ....	17
15. Cederholm, L. (Ed.): Advances in insect systematics and phylogeny. 1981. 415 pp. ....	55
16. Sæther, O. A. Orthoclaadiinae (Diptera: Chironomidae) from the British West Indies, with descriptions of <i>Antillocladius</i> n.gen., <i>Lipurometriocnemus</i> n.gen., <i>Comptosmittia</i> n. gen. and <i>Diplosmittia</i> n.gen. 1981. 46pp. ....	11
17. Ashe, P.: A catalogue of chironomid genera and subgenera of the world including synonyms (Diptera: Chironomidae). 1983. 68 pp. ....	19
18. Enghoff, H.: The milliped genus <i>Cylindroiulus</i> on Madeira — an insular species swarm (Diplopoda, Julida: Julidae). 1982. 142 pp. ....	26
19. Wiederholm, T. (Ed.): Chironomidae of the Holarctic region. Keys and diagnoses. Vol. 1. Larvae. 1983. 457 pp. Hardbound. ....	80
20. Sæther, O. A. & Sublette, J. E.: A review of the genera <i>Doithrix</i> n. gen., <i>Georthocladus</i> Strenzke, <i>Parachaetocladus</i> Wülker and <i>Pseudorthocladus</i> Goetghebuer (Diptera: Chironomidae, Orthoclaadiinae). 1983. 100 pp. ....	20
21. Fjellberg, A.: Arctic Collembola I — Alaskan Collembola of the families Poduridae, Hypogastruridae, Odontellidae, Brachystomellidae and Neanuridae. 1985. 126 pp. ....	20
22. Roth, L. M.: A taxonomic revision of the genus <i>Blattella</i> Caudell (Dictyoptera, Blattaria: Blattellidae). 1985. 221 pp. ....	40
23. Löken, A.: Scandinavian species of the genus <i>Psithyrus</i> Lepeletier (Hymenoptera: Apidae). 1984. 45 pp. ....	10
24. Lomholdt, O.: A reclassification of the larrine tribes with a revision of the Miscophini of southern Africa and Madagascar (Hymenoptera: Sphecidae). 1985. 183 pp. ....	35
25. Nielsen, E. S.: A taxonomic review of the adelid genus <i>Nematopogon</i> Zeller (Lepidoptera: Incurvarioidea). 1985. 66 pp. ....	12
26. Hansson, C.: Taxonomy and biology of the palearctic species of <i>Chrysocharis</i> Förster, 1856 (Hymenoptera: Eulophidae). 1985. 130 pp. ....	20
27. Gärdenfors, U.: Taxonomic and biological revision of Palearctic <i>Ephedrus</i> (Hymenoptera: Braconidae, Aphidiinae). 1986. 95 pp. ....	17
28. Wiederholm T. (Ed.): Chironomidae of the Holarctic region. Keys and diagnoses. Vol. 2. Pupae. 1983. 482 pp. Hardbound. ....	80
29. Sæther, O. A. (Ed.) A conspectus of contemporary studies in Chironomidae (Diptera). Contributions from the IXth Symposium on Chironomidae, Bergen, Norway. 1987. 393 pp. Hardbound. ....	58
30. Cederholm, L. (Ed.): Reports from the Lund University Ceylon Expedition in 1962. Vol. IV. 1988. 160 pp. ....	35
31. Hansson, C. Revision of the New World species of <i>Chrysocharis</i> Förster (Hymenoptera: Eulophidae). 1987. 86 pp. ....	15
32. Bengtson, S.-A. & Enckell, P. H. (Eds.): Island terrestrial invertebrates: Species composition, dispersal and faunal history. 1988. 88 pp. ....	23



33. Ronquist, F. & Nordlander, G. Skeletal morphology of an archaic cynipoid, <i>Ibalia rufipes</i> (Hymenoptera: Ibalidae). 1989. 60 pp. ....	20
34. Wiederholm, T. (Ed.): Chironomidae of the Holarctic region. Keys and diagnoses. Vol. 3. Male imagines. (Diptera). 1989. Hardbound. 538 pp. ....	97
35. Sæther, O. A.: A review of the genus <i>Limnophyes</i> Eaton from the Holarctic and Afrotropical regions (Diptera: Chironomidae, Orthocladinae). 1990. 135 pp. ....	40
36. Scharff, N.: Spiders of the family Linyphiidae from the Uzungwa mountains, Tanzania (Araneae). 1990. 95 pp. ....	40
37. Olafsson, E. Taxonomic revision of wester Palaearctic species of the genera <i>Scatella</i> R.-D. and <i>Lamproscatella</i> Hendel, and studies on their phylogenetic position within the subfamily Ephydrinae (Diptera, Ephydrinae). 1991. 100 pp. ....	45
38. Hansson, C. A catalogue of the Chalcidoidea (Hymenoptera) described by C. G. Thomson, with a checklist of Swedish species. 1991. 70 pp. ....	25
39. Muona, J. A revision of the Indomallesian tribe Galbitini new tribe (Coleoptera, Eucnemidae). 1991. 66 pp. ....	25
40. Enghoff, H. <i>Dolichoiulus</i> — a mostly Macaronesian multitude of millipedes. With a description of a related new genus from Tenerife, Canary Islands (Diplopoda, Julida, Julidae). 1992. 158 pp. ....	55
41. Tjeder, B.† & Hansson, Ch. The Ascalaphidae of the Afrotropical Region (Neuroptera). 1992. 240 pp. ....	70
42. Baranowski, R. Revision of the genus <i>Leiodes</i> Latreille of North and Central America (Coleoptera: Leiodidae). 1993. 149 pp. ....	55
43. Nolte, U. Egg masses of Chironomidae (Diptera). A review, including new observations and a preliminary key. 1993. 75 pp. ....	30
44. Muona, J. Review of the phylogeny, classification and biology of the family Eucnemidae (Coleoptera). 1993. 133 pp. ....	50
45. Pape, Th. The World <i>Blaesoxipha</i> Loew, 1861 (Diptera: Sarcophagidae) 1994. 247 pp. ....	70

We also have small editions of the backnumbers of *Opuscula entomologica* and its supplement series like:

12 Lindroth, C. H. The carabid beetles of Newfoundland including the French islands St. Pierre and Miquelon. 160 pp., 58 figs. ....	18
20. Lindroth, C. H. The Ground-Beetles (Carabidae, excl. Cicindelidae) of Canada and Alaska. Part 2. 1961. 200 pp. ....	18
23. Lindroth, C. H. The faunal history of Newfoundland. Illustrated by Carabid beetles. 1963. 112 pp. ....	10
24. Lindroth, C. H. The Ground-Beetles (Carabidae, excl. Cicindelidae) of Canada and Alaska. Part 3. 1963. 208 pp. ....	18
28. Ball, G. E. A revision of the North American species of the subgenus <i>Cryobius</i> Chaudoir (Pterostichus, Carabidae, Coleoptera). 1966. 166 pp. ....	18
29. Lindroth, C. H. The Ground-Beetles (Carabidae, excl. Cicindelidae) of Canada and Alaska. Part 4. 1966. 240 pp. ....	22
33. Lindroth, C. H. The Ground-Beetles (Carabidae, excl. Cicindelidae) of Canada and Alaska. Part 5. 1968. 296 pp. ....	25
34. Lindroth, C. H. The Ground-Beetles (Carabidae, excl. Cicindelidae) of Canada and Alaska. Part 6. 1969. 248 pp. ....	25
35. Lindroth, C. H. The Ground-Beetles (Carabidae, excl. Cicindelidae) of Canada and Alaska. Part 1. 1969. 48 pp. ....	8

Ask for our complete catalogue!

Freight & check redemption not included. For Swedish deliveries, VAT will be added.

Scandinavian Entomology, Ltd.

Västervång 28, S-247 34 S. Sandby, Sweden

Phone +46/(0)46-518 23, 10 93 30, fax +46/(0)46-5 79 69,

postal giro account 4388668-8.

UNIVERSITÄTSLIBRARIET

95 -09- 2 2

LUND